

College of Nursing and Health Professions



Drexel E-Repository and Archive (iDEA)

<http://idea.library.drexel.edu/>

Drexel University Libraries

www.library.drexel.edu

The following item is made available as a courtesy to scholars by the author(s) and Drexel University Library and may contain materials and content, including computer code and tags, artwork, text, graphics, images, and illustrations (Material) which may be protected by copyright law. Unless otherwise noted, the Material is made available for non profit and educational purposes, such as research, teaching and private study. For these limited purposes, you may reproduce (print, download or make copies) the Material without prior permission. All copies must include any copyright notice originally included with the Material. **You must seek permission from the authors or copyright owners for all uses that are not allowed by fair use and other provisions of the U.S. Copyright Law.** The responsibility for making an independent legal assessment and securing any necessary permission rests with persons desiring to reproduce or use the Material.

Please direct questions to archives@drexel.edu

Preliminary Tests of an Ecological Model of Hispanic Farmworker Health

Abstract

Objective: The purpose of this study was to propose and test an ecological model to structure research and practice concerning farmworker health in the United States. The research question was, “What is the relationship of selected social, cultural, and economic indicators to the health of adult Hispanic migrant farmworkers?”

Design: A model of biogenetic, social, cultural, economic, individual response, and access factors affecting health was derived from the literature and nursing practice. Data from the 1998 National Agricultural Workers Survey (NAWS), conducted by the U.S. Department of Labor, was used in a secondary analysis to perform preliminary tests of the relationships proposed in the model.

Sample: The NAWS conducts interviews with a nationally representative sample of farmworkers employed in the U.S. Because of the importance of theorized cultural factors, the sample for this study was limited to farmworkers who identified themselves as Hispanic, resulting in 1,864 subjects.

Measurements: Variables were used directly from the data or constructed from the available data, and proposed relationships were tested statistically.

Results: Analysis of the data supported the relationships proposed in the model.

Conclusions: The proposed model is a useful tool for organizing variables and giving direction to farmworker health research. Suggestions for future research are made.

Key words: Theoretical model, ecological model, farmworker health

Introduction

The Institute of Medicine (IOM)'s report *Unequal Treatment* makes it clear that disparities in health must be examined from a broad perspective, including socioeconomic status, literacy, and access to health care as well as biology (Smedley, Stith, & Nelson, 2003). This holistic approach is entirely consistent with nursing's long history of viewing patients in the context of their physical and social environments, and the work of public health nurses is rooted in a broad community framework. Unfortunately, this makes the study of "health" in its holistic sense a difficult task, as the public health view implicates multiple potential determinants of health.

The health of non-guest worker farmworkers hired for crop agriculture presents an even greater challenge to researchers, as the social conditions and lifestyle issues surrounding this population are very different from non-farmworker populations. While farmworkers' poverty, low educational levels, and impaired access to health care place them well within the populations included in the IOM report, their seasonal employment, often requiring temporary relocation or migration, cultural issues, and social stigmatization place them apart (Ward & Atav, 2004). Understanding determinants of farmworker health is an important public health issue from the standpoints of social justice, food safety, and infectious disease control. Because of the complexity of farmworkers' lives, however, organizing potential variables can be a daunting task for the public health nurse, whether the intent is to use primary or secondary data for research or to design health-related interventions.

Ecological models have been proposed as a way to structure the consideration of health problems with increasingly complex etiologies (Grzywacz & Fuqua, 2000; Stokols, 2000). These models are based on the ecological perspective that multiple aspects of individuals and

environments interact to influence health (Grzywacz & Fuqua, 2000). Once the complexity of etiologies is understood, interventions can be planned at the individual, family, community, and societal levels (Haughton, 2006; Stokols, 2000).

The purpose of this study was to propose a model of determinants of Hispanic migrant farmworker health, and to test one of the two pathways in the model using data from the National Agricultural Workers Survey (NAWS). The proposed model (see Figure 1) was developed by modifying an explanatory model proposed by Dutch researchers (Uniken Venema, Garretsen, & Van Der Mass, 1995). The Dutch model focused on Surinamese, Turkish, and other immigrant groups who commonly migrate to Holland, understandably assumed a universal health system, and was too complex to be useful in most research studies. The model proposed in Figure 1 aggregated and modified some of the variables from the Dutch model, and added access variables, based on U.S. migrant farmworker health literature (David & Rhee, 1998; Derose & Baker, 2000; Flores, Abreu, Olivar, & Kastner, 1998; Sarver & Baker, 2000); observations and clinical experience of the researcher were also considered in the development of the Determinants of Hispanic Migrant Farmworker Health model.

Variables were grouped into six categories of potential health determinants and arranged into a conceptual model. The major categories included biogenetic variables (age, gender), social, cultural, and economic variables as the primary influences on health. Major intermediate variable categories were individual response concerns and access to health care. Relationships were then theorized among the variables (see Figure 1). The research question was, “What is the relationship of selected social, cultural, and economic indicators to the health of adult Hispanic migrant farmworkers?”

Background

The overwhelming majority of farmworkers today are young, male, and of Mexican origins (Carroll, Samardick, Bernard, Gabbard, & Hernandez, 2005); many who are not Mexican come from other Latin American countries. Despite stricter employment rules established in the 1986 Immigration Reform and Control Act, many farmworkers are undocumented, and some are exploited by labor contractors, who serve as “middle men” between the grower and the laborers (Rothenberg, 1998; Taylor, Martin, & Fix, 1997). Although various farmworker migration patterns have been described, in general some individuals work seasonally in agriculture and do not migrate at all, some migrate annually to work for a single employer and return home at the end of the season, and some “follow the crops”, moving throughout the season to perform specialized work such as harvesting strawberries or tomatoes (Migrant Clinicians Network, 2006). Traditionally, migrant patterns have been described as “streams”, with downstream regions being those states traditionally considered home-base states: California, Texas, and Florida, and upstream regions being those with shorter growing seasons that would provide more temporary employment (Northwest, Midwest, and Northeast). Research indicates that there is a difference in demand and access to health services between home-base and upstream areas (Dever, 1991). These mobility characteristics of farmworkers’ lives place them at increased risk of illness and poor access to health care.

Poor working and housing conditions also disadvantage this population (Early et al., 2006; Mines, Mullenax, & Saca, 2001). The literature indicates that it is difficult to determine working conditions from a survey format; however certain proxies have been developed that are considered relatively accurate (Mines, Mullenax, & Saca, 2001). Among these are payment by piecework rather than hourly or by salary, requiring payment for transportation to work, and

requiring the farmworker to pay for equipment necessary for the job. It should be noted that the proxy of being required to pay for rides to work is not simply having the expense of paying for transportation. It reflects the practice common with some labor contractors of charging a required fee for mandatory labor bus transportation, or requiring workers to pay a third party (*raitero*) for transportation, often standing in the back of a pickup truck. Crowded, inadequate housing is a risk factor for depression (Magana & Hovey, 2003) and makes it difficult to reduce pesticide exposure, thus posing a threat to the physical and mental health and safety of these workers (Early et al., 2006).

Historically, farmworkers have been a marginalized population; since World War II agricultural work in the United States has become “immigrant” work (Massey, Alarcón, Durand, & González, 1987). Language barriers, regulatory restrictions on services available, and low educational levels may further impair access to adequate health care.

Increasingly, migrant farmworkers travel without their families and may be separated from them for months to years (Massey, Durand, & Malone, 2002). It is unclear how much and what kinds of social support these laborers have available during the work season. Studies indicate that the migrant lifestyle is stressful and that farmworkers cope with the stress in both typical and unique ways (Clifford, 1999; Kim-Godwin & Bechtel, 2004; Magana & Hovey, 2003).

Agricultural workers are among the poorest of workers in the U.S. (Runyan, 2000) with mean family incomes of around \$10,000 (Carroll, Samardick, Bernard, Gabbard, & Hernandez, 2005). In addition, for many, migratory employment and/ or undocumented status prevent access to health insurance. Federally funded Community Health Centers provide inexpensive or

free health care to farmworkers, but clinic hours, farm location, poor understanding, and lack of transportation often deter the use of these facilities.

This complexity of context highlights the value of an ecological model when considering farmworker health and its determinants. Preliminary testing of such a model is described in the following section.

Method

Preliminary testing of the proposed model was performed by secondary analysis of data from the National Agricultural Workers Survey (NAWS). The NAWS has the distinction of being one of the largest and most comprehensive sources of data on farmworkers available in the United States. It is a national survey of farmworkers employed in crop agriculture conducted annually since 1989 by the U.S. Department of Labor and made available as a public access data set, with all identifying information removed. The public access data set was analyzed using Statistical Package for Social Sciences (SPSS) 14 for Windows.

Design and Sample

The NAWS uses a complex stratified sampling strategy to interview a representative sample of hired farmworkers in the U.S. (Mehta et al., 2000). Extensive information about demographics, legal status, education, family size and composition, wages, working conditions in farm jobs, and past and planned participation in the U.S. labor force is collected. Only the 1998 data were used for this secondary analysis since specific health questions change from year to year. In addition, the sample for this study was limited to Hispanic respondents since culture was considered an important variable. With these selection criteria, 1,864 interviews were available for analysis.

All 1,864 respondents had complete information on the dependent variable, but some were missing data on correlate variables, notably *family income* with 9% of the sample ($n = 168$) not reporting. Those not reporting an income were younger (27.5 years compared with 33.2 years for those reporting), and more likely to be follow-the crop migrants and undocumented. No attempt was made to impute missing data for these analyses. The logistic regressions (see

analytic strategies) were conducted with the 1672 respondents who had complete data on all variables.

Measures Used

The NAWS contained information relating to biogenetic, social, cultural, economic, and health care access indicators, as well as information on health problems. However, no data on variables from the individual response indicator of the proposed model, such as diet, stress-reduction activities, sleep habits, drug, alcohol, or tobacco use, were available in this survey. Therefore, the model to be tested contained fewer variables than the theoretical model. The individual response indicator was eliminated from the present study and relationships of that indicator were not tested, although it is believed that they exert a major influence on individual health and ability to work and thus they remain in the proposed model. The following section describes the selection, transformation, and integration of NAWS variables into the proposed model.

Variables

Using the model as a guide, the NAWS was reviewed for questions that pertained to the variables of interest. When appropriate, missing or “don’t know” values were set to zero. Some variables used in this study were taken directly from the NAWS data, some were recoded to better represent the concept of interest, and some were constructed by combining portions of several NAWS questions.

Health status.

The dependent variable (*health status*) was operationalized with a single variable that was available directly from the data set (“In the U.S.A., and in the last two years, have you ever had

any health problems (injured or gotten sick)?”). This variable was measured dichotomously (Yes/ No).

Biogenetic indicators

Age and *gender* were available directly in the NAWS data.

Social indicators

Social indicators were operationalized using a number of variables. *Number of years as a farmworker* was available directly from the NAWS data. The literature review indicated that specific *legal status* as available in the NAWS was of less concern than whether the farmworker was documented or undocumented, so a dichotomous variable was created to indicate this differentiation. The discrete *regions* variable available in the NAWS was recoded to reflect whether the farmworker was interviewed upstream or downstream. Initial analysis indicated that farmworkers who “follow the crops” scored differently on many of the variables than either non-migrants or shuttle migrants. The NAWS variable for type of migrant was therefore recoded into a dichotomous variable indicating whether or not the interviewee was a *follow-the-crop* migrant.

The working conditions variables of *payment by piecework* rather than hourly or by salary, requiring payment for transportation to work, and requiring the farmworker to *pay for the equipment* necessary for the job were available in the NAWS but required recoding in order to represent the concepts desired. Two questions in the NAWS asked about transportation payment. One asked if the worker paid someone to take him/her to work, and the other asked if riding the labor bus was mandatory. These were combined to reflect the concept of *payment for riding to work* in a dichotomous fashion.

The survey provided several options of how a worker's pay may be calculated, such as piecework, salary, hourly, or a combination. Since *piecework* or not piecework was the issue of concern, the variable was recoded to reflect this dichotomous concept. The NAWS asked who *paid for equipment* needed by the farmworker; this was recoded to reflect whether the employer paid, or the worker or someone on his/her behalf paid all or some of the equipment cost.

There were many questions in the survey related to farmworkers' *housing* situations, but most were descriptive rather than indicative of quality. One question available directly from the data was how many people, other than family members, the farmworker lived with at the present time, and this was used to operationalize the housing variable.

The most direct measure of educational level, *highest grade completed*, was available directly in the NAWS data. *Literacy* is another measure of education, and one that would account for those who self-educated to some degree. Two NAWS questions about English and primary language literacy were combined, reasoning that literacy in two languages would reduce barriers even more than in only one.

Cultural indicators

Three variables reflective of culture were available in the NAWS data: *English-speaking ability*, *family separation*, and *social support*. English-speaking ability, included in this study to represent language barriers, was available directly from the data, while separation and social support required construction from other variables. The measurement of family separation was available in limited form from questions about how many children less than the age of 18 had been left behind and whether or not a spouse had been left behind for those workers who migrate. Responses to these questions were added, then converted to a dichotomous variable by

recoding to indicate leaving family member(s) at home = 1, not leaving family at home = 0.

Social support, albeit only material support, was measured in a NAWS question regarding sources of material support (church, family, community organizations, charitable organizations, friends) in the past year. These responses were summed to reflect the number of sources of support, then recoded so not having received support = 0, having received support = 1. Making this variable dichotomous lost little in the way of data richness, since few farmworkers had used more than one or two sources of support.

Economic indicators.

Two measures of economic status were used from the NAWS data. The first, *family income*, was available directly and was recorded in fifteen categories ranging from “under \$500” to “over \$40,000.” The second economic measure, *tangible assets*, was constructed from two questions asking if the respondent owned or was buying any of the following in the U.S. or in the home country: plot of land, house, mobile home, car/ truck, business, other. These were added to produce the tangible assets variable with a theoretical maximum of 12, if the farmworker owned all listed assets in both countries.

Access to care indicators.

All three variables used to reflect access to medical care were constructed from other variables available in the data. The barriers variable was constructed from a follow-up to the question “In general, in the U.S.A., would you say that it is easy or difficult for you to get the kind of medical assistance you need?” The follow-up question was “If it is difficult, please

explain why” and was followed by a list of potential barriers. These were added to produce the *barriers* variable used in this study; the potential range was 0 - 11.

Use of government programs targeting the health needs of low-income individuals and families is another variable that could affect health outcomes. The NAWS data included information about use of health-related social programs such as government clinics, Medicaid, and the Women Infants and Children’s (WIC) nutrition program. These were summed, then recoded to reflect no use of government program = 0, use of any of the programs = 1.

Insurance is a variable widely accepted as important to the health of individuals and groups. Employer-provided health and worker’s compensation insurance were available in the NAWS data as three questions addressing free health care for on-the-job illness/ injuries, recuperation, or off-the-job illness/ injuries. The responses to these questions were summed and aggregated into one variable. The variable was then recoded to indicate not having insurance = 0, having any of the forms of insurance = 1. While this variable was intended to address access issues, it also reflects working conditions due to the wording of the questions (“Does your employer provide...?”).

Analytic Strategy

Bivariate analyses were first performed to test the proposed model. Each arrow in the model was considered a hypothesis, and the appropriate statistic was performed to assess the relationship between the variables (data not presented). The relationship of the health problems variable and each correlate variable was also examined with bivariate analysis, either independent samples t-test or Chi square (see Table 2).

The Health Problems variable was then regressed on all correlate variables using logistic regression and a stepwise entry, with the first step encompassing biogenetic, social, and

economic variables, the second step adding cultural variables, and the third step adding access variables (see Table 3). Health problems was also regressed on all correlate variables using the likelihood-ratio test feature of SPSS, in which the model fit is compared successively with and without each variable, thus selecting the set of variables that produce the best model fit (Tabachnick & Fidell, 2007). Significant relationships (at the 0.10 level, see Tabachnick and Fidell, p. 456) from both these regressions, which had most variables in common, were then noted for inclusion in the reduced model.

Finally, of Health Problems was regressed on all variables that were significant in either of the above regressions to produce the reduced model (see Table 4). The large sample ($n = 1672$ with complete data) of the 1998 cycle of the NAWS provided sufficient numbers to conduct this analysis, exceeding the 50 subjects per variable used to estimate the adequacy of sample size (Wright, 1995).

Results

A description of the sample is presented in Table 1. Ages ranged from 14 to 90 years, but on average this was a young (32.7 years) and largely male (84.5%) sample, with over half reporting lack of legal documentation. They reported 0 to 55 years of experience in U.S. farmwork (mean 9.4 years). Educational level and literacy were low, and over 50% reported that they spoke no English at all. Mean family income was \$7,250.

Findings of bivariate analyses indicated that the relationships among indicators were appropriately represented by the proposed model. Specific findings are not the emphasis of this paper, but all relationships hypothesized by the arrows in the tested model were supported by the analysis of these data (not shown). For example, men were more likely than women to lack legal documentation, and undocumented workers had lower family incomes than documented workers. Those with poorer working conditions (paid by piecework or required to pay for rides) also had lower family incomes.

Bivariate analysis of each predictor variable and the dichotomous Health Problems outcome variable indicated a variety of significant findings (see Table 2). Individual biogenetic, social, cultural, economic, and access variables were associated with Health Problems in this population. Older individuals, women, those who had worked longer in U.S. farmwork, and those with low incomes were more likely to report health problems.

When all variables were entered into the stepwise logistic regression, the resulting model was superior to the constant-only model ($p = .000$) and correctly predicted 15% of those with health problems. Significant variables again represented all categories of indicators from the proposed theoretical model (biogenetic, social, cultural, economic, and access). Specifically, gender, follow-the-crop status, location of interview, working conditions as represented by being

required to pay for equipment, literacy, family income, and insurance were significant in this logistic regression model. The non-significant Hosmer and Lemeshow test ($p = .487$) indicated that this model did not differ significantly from the theoretical “perfect” model (Tabachnick & Fidell, 2007). The likelihood ratio test produced a very similar model in which the additional variables of Age and English-speaking ability were identified as contributing to a well-fit model.

Health Problems was regressed on all significant variables from the above models to produce the reduced model. This produced a model that was superior to the constant-only ($p = .000$), not significantly different from the “perfect” model ($p = .582$), and correctly predicted 10.4% of those with health problems. In this analysis, insurance dropped to being non-significant ($p = .093$), but all other variables retained their value as predictors (see Table 4). Thus, the reduced model reflected biogenetic (age, gender), social (follow-the-crop status, region interviewed, working conditions as represented by being required to pay for equipment), economic (family income), cultural (English-speaking ability), and access (insurance, ns) variables.

Discussion

The results of these initial analyses provide some support for use of the Determinants of Hispanic Farmworker Health Model. It is clear that health is a complex concept that responds to numerous determinants, and these analyses indicate that biogenetic, social, cultural, and economic factors are all germane to the study and prediction of health in the Hispanic farmworker population.

Specific findings of this study support many previous reports. For every year of age in this sample, there was a 1% increase in the likelihood of reporting a health problem. Women were almost 2.3 times more likely to report health problems than were men, possibly because of the physically demanding work and possibly because they were more likely to admit to difficulties (Mines et al., 2001). Follow-the crop migrants and those interviewed upstream were more likely to report health problems, a finding that supports the earlier work of Dever (1991). Not surprisingly, poor working conditions were also associated with a higher likelihood of reporting health problems in this population, as had been previously reported for a California-based sample (Mines et al., 2001).

Three seemingly counter-intuitive findings were that better English speaking ability, higher family incomes, and having insurance were associated with poorer reported health. It is possible that facility in the local language increased the respondents' comfort level in responding affirmatively to the question, or that the effect of acculturation (often measured by language spoken) resulted in a different interpretation of the question. Higher incomes have been associated with poorer self-rated health worldwide (Sadana, Mathers, Lopez, Murray, & Iburg, 2002), possibly related to higher expectations. This phenomenon may partially explain the income finding as well as the insurance finding of the present study. In this population, having

insurance and having a higher income may result in a level of confidence which permitted these workers to admit having had health problems.

For the purposes of testing the model, however, the more important issue was that a wide variety of variables were significant in the prediction of health problems. No single predictor was predominant, and the model that included all 22 variables correctly predicted 15% of health problems, while the more parsimonious reduced model predicted 10%. These findings support an ecological approach to explaining, predicting, and studying health. The model can be used, as it was in this study, to organize data from a large dataset, or it could lend structure to smaller, more focused research. The findings from this study indicate that an ecological approach to migrant farmworker health is valid.

This study reinforces what public health nurses who work with migrant farmworkers have always known: more than health service availability is required to improve this population's health. Nurses working in "upstream" areas and those who see follow-the-crop migrants can request funding for more outreach and case-finding services, especially for groups with poorer working conditions. Culturally appropriate health services for female farmworkers are especially needed. This study also lends support for nurses to be politically active in improving housing and working conditions for this essential workforce.

Limitations to this study stem largely from the fact that it was a secondary analysis and therefore constrained to the variables available in the data set which, while extensive, were sometimes not ideal. Absence of variables pertaining to individual response, where most nursing interventions would be targeted, and the nature of the "health problems" variable are most notable as limitations (Ward, 2007). In addition, the data were cross-sectional and therefore causal relationships could not be inferred. An area not addressed in these analyses but clearly

critical to farmworkers' health is that of physical environment, including exposure to pesticides and exposure to the elements while working. This could be considered as a future addition to the Determinants of Migrant Farmworker Health Model.

Additional research is needed in almost all areas of farmworker health. For researchers with appropriate language and cultural skills, qualitative studies inquiring what it is like to follow the crops while managing acute or chronic conditions would help health care providers deliver meaningful, appropriate care. Additional inquiry could illustrate if male and female farmworkers' health concerns are the same, and how they decide when to seek health care. Perhaps the most pressing need for further research, and critically important to the public health nurse, is the testing of the individual response category. Understanding the status of farmworkers' diets, health behaviors, responses to stress, and sleep would be helpful in guiding nursing actions both in the clinical and policy arenas. This model could also be tested for use with other culturally distinct groups, and could be especially effective if Health Related Quality of Life were the outcome variable.

Multiple factors are involved in the etiology of health problems for Latino farmworkers in the United States, and the health of farmworkers is of critical importance to the public. The findings of this study provide initial support for interventions at the local, community, and policy levels to modify conditions associated with poorer health in this population. Examining health from this broad perspective provides Public Health Nurses with a framework for both research and interventions with Latino farmworkers.

References

- Carroll, D., Samardick, R. M., Bernard, S., Gabbard, S., & Hernandez, T. (2005). *Findings from the National Agricultural Workers Survey (NAWS), 2001 - 2002: A Demographic and Employment Profile of United States Farmworkers* (No. Research report # 9). Washington, DC.
- Clifford, S. (1999). Addressing emotional health among Latino farmworkers: A North Carolina Experience. *MCN Streamline*, 5(5), 1-3.
- David, R. A., & Rhee, M. (1998). The impact of language as a barrier to effective health care in an underserved urban Hispanic community. *The Mount Sinai Journal of Medicine*, 65(5&6), 393-397.
- Derose, K. P., & Baker, D. W. (2000). Language difficulties are as much of a barrier to health care for some Latinos as being uninsured. *AHRQ*, 237(May).
- Dever, G. E. A. (1991). *Migrant health status: Profile of a population with complex health problems*. Austin, Texas: Migrant Clinicians' Network Monograph Series.
- Early, J., Davis, S. W., Quandt, S. A., Rao, P., Snively, B. M., & Arcury, T. A. (2006). Housing characteristics of farmworker families in North Carolina. *Journal of Immigrant and Minority Health*, 8(2), 173 - 184.
- Flores, G., Abreu, M., Olivar, M. A., & Kastner, B. (1998). Access barriers to health care for Latino children. *Archives of Pediatric and Adolescent Medicine*, 152, 1119-1125.
- Grzywacz, J. G., & Fuqua, J. (2000). The social ecology of health: Leverage points and linkages. *Behavioral Medicine*, 26, 101 - 115.

- Haughton, B. (2006). Applying the social-ecological model to nutrition issues that promote health and prevent disease. *Family & Community Health, 29*(1), 3 - 4.
- Kim-Godwin, Y. S., & Bechtel, G. A. (2004). Stress among migrant and seasonal farmworkers in rural southeast North Carolina. *Journal of Rural Health, 20*(3), 271-278.
- Magana, C. G., & Hovey, J. D. (2003). Psychosocial stressors associated with Mexican migrant farmworkers in the midwest United States. *Journal of Immigrant Health, 5*(2), 75-86.
- Massey, D., Alarcón, R., Durand, J., & González, H. (1987). *Return to Aztlan: The Social Process of International Migration from Western Mexico*. Berkely, CA: University of California Press.
- Massey, D. S., Durand, J., & Malone, N. J. (2002). *Beyond smoke and mirrors: Mexican immigration in an era of economic integration*. New York: Russell Sage Foundation.
- Mehta, K., Gabbard, S., Barrat, V., Lewis, M., Carroll, D., & Mines, R. (2000, 3/28/2000). Findings from the National Agricultural Workers Survey (NAWS) 1997-8: A demographic and employment profile of United States farmworkers. Retrieved 9/24, 2000, from <http://www.dol.gov/dol/asp/public/programs/agworker/naws.htm>
- Migrant Clinicians Network. (2006). Migration Patterns. Retrieved August 15, 2006, from http://www.migrantclinician.org/migrant_info/migration_patterns.php
- Mines, R., Mullenax, N., & Saca, L. (2001). *The Binational Farmworker Health Survey: An In-depth Study of Agricultural Worker Health in Mexico and the United States* (Research Report). Davis, California: California Institute for Rural Studies.
- Rothenberg, D. (1998). *With these hands: The hidden world of migrant farmworkers*. New York: Harcourt Brace.

- Runyan, J. L. (2000). *Profile of Hired Farmworkers, 1998 Annual Averages* (Agricultural Economic Report No. 790). Washington, D.C.: U.S. Department of Agriculture.
- Sadana, R., Mathers, C. D., Lopez, A. D., Murray, C. J., & Iburg, K. (2002). Comparative analysis of more than 50 household surveys on health status. In C. J. L. Murray, J. A. Salomon, C. D. Mathers & A. D. Lopez (Eds.), *Summary measures of population health: Concepts, ethics, measurement, and applications*. Geneva: World Health Organization.
- Sarver, J., & Baker, D. W. (2000). Effect of language barriers on follow-up appointments after an emergency department visit. *Journal of General Internal Medicine*, 15(4), 256-264.
- Smedley, B. D., Stith, A. Y., & Nelson, A. R. (Eds.). (2003). *Unequal Treatment: Confronting Racial and Ethnic Disparities in Health Care*. Washington, D.C.: The National Academies Press.
- Stokols, D. (2000). Social ecology and behavioral medicine: Implications for training, practice, and policy. *Behavioral Medicine*, 26, 129 - 138.
- Tabachnick, B.G., & Fidell, L.S. (2007). *Using multivariate statistics, 5th ed.* Boston: Pearson Education.
- Taylor, J. E., Martin, P. L., & Fix, M. (1997). *Poverty amid prosperity: Immigration and the changing face of rural California*. Washington, D.C.: The Urban Institute Press.
- Uniken Venema, H. P., Garretsen, H. F. L., & Van Der Mass, P. J. (1995). Health of migrants and migrant health policy: The Netherlands as an example. *Social Science & Medicine*, 41(6), 809-818.
- Ward, L. S. (2007). Farmworker health status: A review of measurement methods. *Hispanic Health Care International*, 5(1), 34 - 40.

- Ward, L. S., & Atav, A. S. (2004). Migrant Farmworkers. In N. Glasgow, L. Morton & N. Johnson (Eds.), *Critical Issues in Rural Health* (pp. 169 - 181). Ames, Iowa: Blackwell Publishing.
- Wright, R.E. (1995). Logistic Regression. In L.G. Grim & P.R. Yarnold (Eds.), *Reading and understanding multivariate statistics*. Washington, D.C.: American Psychological Association.

Table 1. Description of the Hispanic Farmworker Sample on Selected Characteristics

Characteristic	n	%	Mean (s.d.)
Health Problems: Yes	472	25.4	
No	1388	74.6	
Biogenetic			
Age (mean)	1857		32.7 (12.0)
Gender: Male	1572	84.5	
Female	288	15.5	
Social			
Legal status: Documented	862	46.9	
Undocumented	976	53.1	
Follow-the-crop	429	23	
Not follow-the-crop	1435	77	
Region: Downstream	1166	62.6	
Upstream	698	37.4	
Piecework: No	1464	78.5	
Yes	400	21.5	
Pay for ride: No	1424	76.4	
Yes	440	23.6	
Pay for equipment: No	1371	73.6	
Yes	479	25.7	
Years in farmwork	1862		9.36 (9.24)

Characteristic	n	%	Mean (s.d.)
# living with	1853		3.85 (4.32)
Highest grade	1854		6 (3.41)
Literacy level (0 – 6 potential)	1835		2.92 (1.43)
English literacy: Not at all	1134	60.8	
A little	470	25.2	
Somewhat	114	6.1	
Well	144	7.7	
Native language literacy: Not at all	129	6.9	
A little	209	11.2	
Somewhat	418	22.4	
Well	1081	58.0	
Cultural			
English speaking ability: None	942	50.5	
A little	607	32.6	
Somewhat	145	7.8	
Well	169	9.1	
Family separation: No	1355	72.7	
Yes	509	27.3	
Social support: No	1555	83.4	
Yes	309	16.6	
Economic			
Family income	1696		\$7250

Characteristic	n	%	Mean (s.d.)
Tangible assets	1692		1.2 (0.85)
Access			
Barriers:	1582		0.75 (1.0)
Used gov't programs: No	1546	84	
Yes	302	16	
Insurance: No	585	68.6	
Yes	1279	31.4	

Table 2. Bivariate Relationships of Selected Characteristics and Health Problems (mean values or percent)

Characteristic		Health Problems n = 472	No Health Problems n = 1388	Sig.	CI 95% for mean values
Age		m = 35	32	.000	(-3.99, -1.49)
Gender:	Male	22.3%	77.7%	.000	
	Female	42.0%	58.0%		
Legal status:	Documented	31.2%	68.8%	.000	
	Undocumented	19.6%	80.4%		
Follow-the-crop:	No	24.7%	75.3%	.188	
	Yes	27.8%	72.2%		
Region:	Downstream	23.6%	76.4%	.020	
	Upstream	28.4%	71.6%		
Paid by piecework:	No	25.5%	74.5%	.871	
	Yes	25.1%	74.9%		
Pay for ride:	No	25.6%	74.4%	.669	
	Yes	24.6%	75.4%		
Pay for equipment:	No	23.8%	76.2%	.012	
	Yes	29.6%	70.4%		
Years in farmwork		m = 10.89	8.85	.000	(-3.00, -1.08)
# living with (housing)		m = 3	4	.000	(0.47, 1.37)
Highest grade		m = 6.22	6.04	.333	(-0.54, 0.18)

Characterstic	Health Problems n = 472	No Health Problems n = 1388	Sig.	CI 95% for mean values
Literacy level (7 categories)	m = 3.16	2.83	.000	(-0.47, -0.17)
English speaking ability: None	19.6%	80.4%		
A little	28.7%	71.3%	.000	
Somewhat	35.4%	64.6%		
Well	37.3%	62.7%		
Family separation: No	26.4%	73.6%	.102	
Yes	22.7%	77.3%		
Social support: No	26.5%	73.5%	.013	
Yes	19.7%	80.3%		
Income (15 categories)	m = 6.90	5.58	.000	(-1.64, -1.00)
Tangible assets	m = 1.32	1.16	.001	(-0.25, -0.07)
Barriers to health care	m = .73	.75	.616	(-0.08, 0.14)
Use of gov't programs: No	23.1%	76.9%	.000	
Yes	37.1%	62.9%		
Insurance: No	26.8%	73.2%	.042	
Yes	22.3%	77.7%		

Table 3. Odds Ratios (95% CI) of Health Problems, Stepwise Regression Analysis (n = 1199)

Characteristic	Odds ratio (95% CI)		
	Step 1	Step 2	Step 3
Age	1.01 (0.99-1.03)	1.01 (1.00-1.03)	1.01 (1.00-1.03)
Female gender	2.39 (1.70-3.34)	2.36 (1.67-3.35)	2.38 (1.67-3.38)
Undocumented	0.87 (0.61-1.24)	0.89 (0.62-1.27)	0.91 (0.63-1.31)
Follow-the-crop	1.51 (1.09-2.09)	1.55 (1.12-2.15)	1.59 (1.15-2.14)
Interviewed upstream	1.43 (1.07-1.91)	1.42 (1.06-1.89)	1.49 (1.11-2.00)
Paid by piecework	0.85 (0.60-1.19)	0.86 (0.61-1.22)	0.88 (0.62-1.25)
Pay for transport	1.20 (0.88-1.63)	1.23 (0.90-1.68)	1.21 (0.88-1.66)
Pay for equipment	1.31 (0.97-1.77)	1.30 (0.96-1.76)	1.35 (0.99-1.84)
Years in farmwork	1.00 (0.98-1.02)	1.00 (0.98-1.02)	1.00 (0.98-1.02)
More crowded housing	1.01 (0.98-1.05)	1.02 (0.98-1.05)	1.01 (0.98-1.05)
Highest grade	1.00 (0.95-1.05)	1.00 (0.95-1.05)	1.00 (0.95-1.05)
Better combined literacy	1.13 (0.99-1.28)	1.05 (0.91-1.22)	1.07 (0.92-1.24)
Higher family income	1.10 (1.04-1.16)	1.09 (1.03-1.15)	1.08 (1.01-1.14)
More tangible assets	1.08 (0.92-1.27)	1.10 (0.93-1.30)	1.09 (0.92-1.29)
Better English-speaking ability		1.19 (0.96-1.47)	1.19 (0.96-1.48)
Has social support		1.09 (0.70-1.70)	1.06 (0.67-1.65)
Family separation		0.89 (0.64-1.23)	0.91 (0.65-1.27)
Barriers to health care			1.09 (0.95-1.24)
Had Insurance			1.46 (1.06-2.01)

Characteristic	Odds ratio (95% CI)		
	Step 1	Step 2	Step 3
Used government programs			1.20 (0.85-1.69)
Block Chi-square (sig.)	76.56 (0.000)	3.29 (0.349)	7.77 (0.051)
% with Health Problems correctly predicted	11.0	12.5	15.0

Table 4. Reduced Model for Logistic Regression of Health Problems on Selected Characteristics

Characteristic	OR	95% CI
Age	1.01	(1.01, 1.02)
Female gender	2.27	(1.70, 3.03)
Follow-the-crop	1.47	(1.12, 1.95)
Interviewed upstream	1.46	(1.15, 1.86)
Required to pay for equipment	1.43	(1.10, 1.85)
Higher family income	1.13	(1.08, 1.18)
Better English-speaking ability	1.22	(1.08, 1.38)
Had insurance	1.27	(0.96, 1.67)

Model Chi Square p = 0.00

Hosmer and Lemeshow goodness-of-fit: p = .582

Predicted Health Problems: 10.4%

Figure 1: Proposed Theoretical Model of Determinants of Hispanic Migrant Farmworker Health

